Missouri Spring Rise – Spring-Rise Alternative

Hydrology/Water Quality Technical Group

Missouri River Plenary Meeting

July 26, 2005



a. Number of Rises

One/Two depending on storage conditions

- No Spring Rise below 31.0 MAF storage on March 15
- A single rise between 31.0 and 34.0 MAF
- Two rises between 34.0 and 58.5 MAF



b. Flood Control Targets/Constraints

Modify the flood control constraints minimally during the spring rise period.



c. First Rise

i. Timing

Coincide first rise with the start of navigation support releases from Gavins Point



c. First Rise

ii. Duration and rise and fall rates

- Rise at a rate of 6,000 cfs/day
- 2 day peak
- Fall 4,000 first day, then prorate the fall rate so that the total length of the rise is 16 days



c. First Rise

iii. Magnitude

- Prorate between 22,000 cfs and 6,000 cfs
- Use an absolute flow cap of 35,000 cfs releases from Gavins Point



d. Flows between rises

Use Master Manual curves

 Use Flow to Target except during evacuation periods



e. Second Rise

- No rise when system storage is below 31.0 MAF
- Prorate the rise between 31.0 MAF and 54.5 MAF
- Above 58.5 MAF have no rise



e. Second Rise

i. Timing

 As late as possible – must consider the bird species and avoid unacceptable levels of take



e. Second Rise

ii. Duration and rise and fall rates

- Rise up to 6,000 cfs/day
- 2 day peak
- Drop 4,000 cfs/day for two days, prorate the fall rate so that the total length of the rise is 21 to 28 days



e. Second Rise

iii. Magnitude

- Prorate between 20,000 cfs and 10,000 cfs
- Use an absolute flow cap of 48,000 cfs releases from Gavins Point



f. How to address water availability and variation for wet, normal, and dry years

- Use March 15th storage check to set number of peaks and their magnitudes.
- Based on navigation support and drought conservation in the Current Water Control Master Manual.



g. Volume of Water Used

Uses an additional 0.34 MAF compared to Current Water Control Plan




h. Proposal Flexibility

- The Corps should use all forecasting abilities to reduce flooding
- The Corps should be able to react to unexpected events during the spring rise period
- As more tern and plover habitat is created, the spring rise may be shifted later in June
- Proposed flow rates could be targeted immediately below the James River confluence.


Release Hydrograph

Normal System Storage - Bimodal Rise




Release Hydrograph

Low System Storage - Bimodal Rise



Release Hydrograph

Low System Storage - Single Rise



Questions?